

#### NC Department of Health and Human Services NC Nurse Aide I Curriculum

Module AA Measurement

July 2024

1

#### **Objectives**

- Define vital signs and measurement skills nurse aides perform
- 2. List equipment needed to measure:
  - Temperature
  - Pulse
  - Respirations
  - · Blood pressure
  - Height
  - Weight
  - · Intake and output of fluids

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

2

#### **Objectives**

- Compare and contrast the various thermometers used to measure temperature
- 4. Label components of equipment used to measure blood pressure, height, and weight
- 5. Identify normal findings for temperature, pulse, respirations, and blood pressure
- 6. Locate sites used to measure temperature, pulse, respirations, and blood pressure

#### **Objectives**

- Explain how to use equipment that measures temperature, pulse, respirations, blood pressure, height, weight, intake of fluids and output.
- Differentiate units of measurement nurse aides use during the care of residents – millimeters of mercury (mmHg), degrees Fahrenheit, ounces, milliliters (mL), cubic centimeters (cc), inches, and feet.
- 9. Convert ounces to milliliters and inches to feet/inches.

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

4

#### **Vital Signs**

- Measurable variables that indicate the general state or condition of a person: body temperature, pulse rate, respiratory rate, and blood pressure
- Also called TPR & BP

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

5

#### 5th Vital Sign - Pain

- Pain is often considered the 5<sup>th</sup> vital sign and is as important as other vital signs
- Pain is subjective and is whatever the patient says it is and response varies from patient to patient
- A nurse aide uses a pain scale to identify pain levels and then reports to the nurse
- Refer to Module V for more in-depth information



#### Why Check Vital Signs?

- Changes can indicate worsening of a resident's condition
- Can reflect response to medication and/or treatment
- Value of a vital sign may be basis for a medication or change of a medication
- · Accuracy when taking vital signs is crucial
- Report abnormal vital signs immediately to nurse and per facility policy

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

7



8


# What Is an Analog Watch and Why Is it Included in EQUIPMENT for Vital Signs?

- A watch that has moving hands and is typically marked with numbers 1 – 12
- It has an hour hand, minute hand, and second hand
  - \*\*IMPORTANT\*\*
- Nurse aides use the second hand to count respirations and pulse rate

	Con the	12	J.	Hand
Hour	210		2*	
land	.8		4.	Second
	1	6 5	Service .	

# Analog Watch – How to Count for Full 60 Seconds for Respirations and Pulse

- First, identify what number the second hand is on
- Second, while watching the second hand, start counting the vital sign you want to know and stop counting on the same number when secondhand returns to that number
- We will practice once we learn how to check for a pulse and respirations



10

#### **Body Temperature**





Beneficial to identifying issues of the immune system and response to medical treatment.

- · Common terminology
  - Fever elevated temperature
  - Febrile with a fever
  - Afebrile without a fever
  - Thermometer device to measure body temperature
  - Fahrenheit and Centigrade

11

### Factors Affecting Temperature

- Age
- Illness
- Stress
- Environment
- Exercise
- Time of day





#### **Sites for Checking Temperature**

- · Mouth (oral)
- · Rectum (rectal)
- · Armpit (axilla)
- Ear (tympanic)
- · Temporal (forehead)



Important- Check with nurse or care plan to see what type of thermometer should be used

13

#### Types of Thermometers

- Digital oral, rectal, axillary
- Electronic oral, rectal, axillary
- Tympanic ear
- Temporal forehead
- Non-mercury, liquid-filled glass (oral green tipped; rectal red tipped)



14

#### **Temperature Values**

- Oral
  - Baseline 98.6°F
  - Normal range 97.6°F to 99.6°F
- Rectal
  - Baseline 99.6°F
  - Normal range 98.6°F to 100.6°F
- Axillary
  - Baseline 97.6°F
  - Normal range 96.6°F to 98.6°F



4	
	-

#### **Temperature Values**

- Tympanic membrane
  - Baseline 98.6°F
  - Normal range 97.6∘F to 99.6∘F
- Temporal
  - Baseline 98.6°F
  - Normal range 97.6°F to 99.6°F



16

#### **Digital Thermometer**

- Oral, rectal, or axillary
- Displays results digitally within 2 60 seconds; beeps or flashes when done
- · Battery-operated
- Requires a disposable sheath



17

#### **Electronic Thermometer**

- · Oral, rectal, or axillary
- Has oral/axillary and rectal probes; requires probe cover
- $\bullet$  Displays results digitally 2  $-\,60$  seconds; beeps or flashes when done
- Battery-operated; stored in recharging device
- Requires a probe cover



#### **Tympanic Thermometer**

- Ear
- Registers temperature in seconds



19

#### **Temporal Thermometer**


- Measures heat from skin over the forehead, specifically over temporal artery
- Done by a stroke or scan over the area
- Registers within 3 seconds
- Noninvasive


20


#### Non-mercury, Liquid-filled Glass Thermometer

- · Oral, rectal, or axillary
- Color-coded; blue or green for oral; red for rectal
- Takes a longer time to register
- Nurse aide must read at eye level after it registers temperature; held at stem
- Most use Fahrenheit scale to measure temperature









#### When NOT to Take an Oral Temperature

- Unconscious
- Recent facial/mouth surgery
- Recent injury to face
- Sores/redness/mouth pain



- Confused/agitated
- History of seizure
- Using oxygen
- Mouth-breather
- Feeding tube

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024 25



- · Has diarrhea
- · Has rectal problem
- · Has heart disease
- Recent rectal surgery
- · Is confused or agitated



NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

26

#### Pulse

The beat of the heart felt at an artery, as a wave of blood passes through the artery

- · Pulse rate
- · Pulse rhythm
- · Pulse force



# Pulse Sites Temporal Carotid Apical Brachial Radial Pedal Popilical Popilical

28

# Radial Pulse Site Typically used to take pulse during routine vital signs checks Does not expose resident Located on thumb side of wrist Nurse aide places their first 2 or 3 fingers (never thumb) over the radial pulse

29

# Pulse Values • Normal between 60 and 100 beats per minute • Regular • Strong • Refer to Module H for abnormalities

30

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

# Counting Pulse – Equipment • Analog watch with second hand • Notepad and pen

31

#### **Checking Pulse and Documentation**

- Nurse aide counts pulse rate for 60 seconds
- While watching second hand of watch, the nurse aide starts and stops counting on same number
- · Document on record if pulse is normal
- Document on record and notify nurse if pulse is abnormal

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

32

#### Checking Pulse - Example #1

- Nurse aide begins counting pulse rate when second hand is on 4 and stops counting pulse rate when the secondhand lands on
- Nurse aide counts a pulse rate of 82 in 60 seconds
- 82 is the number the nurse aide would document



#### Checking Pulse - Example #2

- Nurse aide begins counting pulse rate when the second hand is on 10 and stops counting pulse rate when the second hand lands on \_\_\_\_\_
- Nurse aide counts a pulse rate of 109 in 60 seconds
- 109 is the number the nurse aide would document
- Nurse aide would notify the nurse; why?



NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

34

#### Respiration

- Process that supplies oxygen to cells and removes carbon dioxide from cells
- · Involves
  - Inspiration (inhalation) breathing in oxygen; chest rises
  - Expiration (exhalation) breathing out carbon dioxide; chest falls
- Each respiration = 1 inspiration and 1 expiration
- Respiratory rate (or respirations) the number of inspirations/exhalations a person takes in a minute

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

35

#### **Respiration Values**

- · Normal breathing is defined as eupnea
- · Between 12 and 20 breaths/minute
- Regular
- Quiet
- · Both sides of chest equal
- · For abnormalities refer to Module H



#### **Counting Respirations – Equipment**

- · Analog watch with second hand
- · Notepad and pen



37

#### **Observation and Documentation**

- Nurse aide counts respirations for 60 seconds
- 1 respiration = 1 inspiration (chest rising) and 1 expiration (chest falling)

Inspiration and Expira



- While watching second hand of watch, nurse aide starts and stops counting on same number
- Document on record if respirations are normal
- Document on record and notify nurse if respirations are abnormal

38

#### **Stealth Respirations**

- Merriam Webster Dictionary defines stealth as "a cautious, unobtrusive, and secretive way of moving or proceeding intended to avoid detection"
- Check respirations right after checking pulse (without moving hand from wrist)
- Why? The patient doesn't change their breathing pattern, and the results are more accurate when they do not know their breathing pattern is being observed

#### Checking Respirations - Example #1

- Nurse aide begins counting respirations when the second hand is on 4 and stops counting respirations when the secondhand lands on \_\_\_\_\_
- Nurse aide counts 16 chest rises/falls in 60 seconds
- 16 is the number the nurse aide would document



40

#### Checking Respirations - Example #2

- Nurse aide begins counting respirations when the second hand is on 10 and stops counting respirations when the second hand lands on \_\_\_\_\_
- Nurse aide counts 24 chest rises/falls in 60 seconds
- 24 is the number the nurse aide would document
- Nurse aide would notify the nurse; why?



41

#### **Blood Pressure**




- The amount of force exerted by the blood against the walls of the artery
- Top number is the systolic pressure; the pressure when the heart contracts and pumps blood out
- Bottom number is the diastolic pressure; when the heart rests as the heart fills with blood


#### **Blood Pressure**

- Important indicator of health status; shows how well the heart is working
- Can change from minute to minute depending on:
  - -Activity of the resident
  - -Lifestyle choices
  - -Reaction to stress
  - -Acute injury or emergency
  - -Medications

43



44



#### The Blood Pressure Value

- Measured in millimeters of mercury (mmHg)
- · Recorded as a fraction
- Pronounced as 120 over 80

120 (systolic) 80 (diastolic)

46

#### **Normal Blood Pressure Ranges**

American Heart Association defines normal blood pressure as less than 120/80

- Systolic Less than 120 mm Hg
- Diastolic Less than 80 mm Hg
- · Document on record
- If above 120/80, document on the record and notify the nurse
- Refer to Module H for abnormalities

47

#### **Using What You Have Learned**

Evaluate the health of individuals on the next 3 slides



#### Example #1

- BP = 116/72
- · A 64-year-old female
- Weighs 130 pounds
- She has finished baking an apple pie
- She doesn't smoke or drink



49

#### Example #2

- BP = 162/86
- A 72-year-old male
- Weighs 260 pounds



- He just ate a couple of hot dogs and potato chips for lunch and is on his 4th beer
- He just sent his son to the store for more cigarettes



50

#### Example #3

- BP = 180/94
- A 22-year-old male
- Weighs 170 pounds





- He just wrecked his brand new truck
- · He has a broken leg
- He is on a stretcher in the back of an ambulance

#### **Checking Blood Pressure**



The nurse aide uses 3 senses when checking a blood pressure

- Seeing watches needle's movement in relation to numbers on the manometer
- Hearing uses the stethoscope to listen for changes in blood flow in the brachial artery
- Touching controls inflation and deflation of cuff using thumb and index finger

52

#### **Checking Blood Pressure - Equipment**

- Stethoscope
- Sphygmomanometer, also known as blood pressure cuff (BP cuff)
- · Alcohol wipes
- · Notepad and pen




53

#### Stethoscope



- · Instrument used to listen to heart and lung sounds
- For blood pressure checks, used to listen to sounds in brachial artery
- May be single-head (with diaphragm only) or dualhead (with diaphragm and bell)



55

#### Stethoscope - Ear Pieces

- Always clean before and after use with alcohol wipes
- Insert ear pieces into ears so they point forward toward the nose
- Should fit snugly in ears




56

#### **Dual-head Stethoscope – Diaphragm**

Before using dual-head stethoscope to take blood pressure, determine which side of chest-piece is active; to check blood pressure, diaphragm needs to be active






58

#### **Diaphragm Concepts**

- · Always clean diaphragm with alcohol wipes
- Warm diaphragm before making contact with resident
- To use diaphragm, apply enough pressure to make seal against brachial artery



59




#### Electronic (Digital)

- No stethoscope needed
- After BP cuff is placed on arm, button is pressed causing cuff to inflate/deflate automatically
- BP reading is displayed



61





62

# Manometer

- Marked with long and short lines and has a needle
- Long lines mark 10 mm Hg
- Short lines mark 2 mm Hg







# Example of Blood Pressure Reading • Systolic is 150 mm Hg • Diastolic is 88 mm Hg • BP is written down as 150/88 • BP is pronounced as "150 over 88" Systolic is 150 mm Hg Diastolic is 150 mm Hg Diastolic is 150 mm Hg Diastolic is 150 mm Hg NCDHHS/IPHSR/HCPEC | Module AA Measurement | July 2024

#### Cuff

- After wrapping the cuff around the bare upper arm
- The cuff inflates and puts pressure on the brachial artery
- As cuff deflates, BP is determined



NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

67

#### Cuff

- Cuff sizes come in child, small, regular, and extra-large
- Important to choose correct size



NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

68

#### Cuff

- Typically has 1 or 2 arrows (left arm/right arm) on cuff which align with brachial artery
- Cuff positioned/wrapped at least an inch above the elbow
- Cuff or stethoscope should not be placed over clothing



NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

#### **Tubing**

- Made of rubber
- Two tubes connect the:
   (#1) Cuff to the manometer and
   (#2) Cuff to the handheld inflation bulb



NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

70

#### Inflation Bulb with Air-release Valve

To inflate cuff, turn air-release valve clockwise to close valve; then squeeze the bulb; remember thumb goes up, needle goes up





NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

71

#### Inflation Bulb with Air-release Valve

To deflate cuff and open valve, turn air-release valve counterclockwise with the thumb and index finger in a slow and controlled manner; remember thumb goes down, needle goes down





NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

#### Inflation Bulb with Air-release Valve

- Inflate cuff to between 160 mm Hg to 180 mmHg
- If beat is heard immediately, deflate the cuff; wait 30 60 seconds; inflate cuff to no more than 200 mmHg



73

#### Inflation Bulb with Air-release Valve

Goal: Learn how to inflate the cuff and how to deflate the cuff in a slow, controlled manner



NCDHHS/DHSR/HCPEC | Module AA Measurementi July 2024

74

#### **Blood Pressure - Tips During Procedure**

- Do not take blood pressure on an arm with an IV, dialysis shunt, or another medical device
- Avoid taking blood pressure on a side that has been injured, burned, paralyzed, has a cast, or has had a mastectomy



- Do not place the blood pressure cuff over clothing
- Make sure resident has an empty bladder
- · Limit conversation
- Support the arm at heart level
- Make sure back is supported
- Make sure legs are not crossed

#### **Checking for Orthostatic Hypotension**

- Blood pressure is abnormally low
- BP checked while lying down, record on notepad



- Have resident sit up, wait 2 minutes, check BP, record on notepad
- Have resident stand up, wait 2 minutes, check BP, record on notepad
- Throughout process, check to see if resident is feeling weak, dizzy, faint, or seeing spots
- · Record and report to nurse

76

#### Height and Weight- Overview

- Height and weight measured on admission to the facility
- Units of measure per facility policy
  - Height may be measured using feet and inches or just inches
  - Weight may be measured in pounds or kilograms
- · After admission
  - Height typically not measured again
  - Weight measured per facility policy and/or doctor's orders as directed by nurse and care plan – daily, weekly, monthly


NCDHHS/DHSR/HCPEC | Module AA Measurement | July 202

77

#### Physician Mechanical Beam Scale

- Used for measuring height and weight
- For residents who cannot stand, measure
  - Height in bed using tape measure and ruler
  - -Weight using a chair, wheelchair, bed, or mechanical lift, as directed by nurse or care plan





79

#### **Height Component – Head Piece**


- Becomes active when extended upward in preparation for measuring resident's height
- Lowered and placed on resident's head and height measured
- Becomes at rest when flat and low against height rod




80

# Height Component – Rod • 2 Units of measure - Inches - Centimeters • 2 sections - Movable upper section - Non-movable lower section

# Height Component - Rod • Movable upper section - Raised or lowered to adjust to resident's height - "Read height here" area is the location of the height value if resident's height is located in this section - Numbers increase from top to bottom • Non-movable lower section - Height read in lower section if resident's height is located in this area - Numbers increase from bottom to top





#### Converting Inches into Feet/Inches

- To convert inches to feet and inches
  - Divide the number of inches by 12
  - The quotient is the feet and the remainder of inches

1 foot = 12 inches

85

#### Height in Feet and Inches

- Resident's height is 68 inches
- How does the nurse aide convert 68 inches to feet and inches?
  - 1 foot = 12 inches
  - Divide 68 inches by 12 inches
  - Quotient = 5, remainder = 8
- Answer = 5 feet, 8 inches



Resident's height is 68 inches or 5 feet, 8 inches

NCDHHS/DHSR/HCPEC | Module AA Measurement | July 2024

86

#### Weights of Residents in Long-term Care

- In general, standards of practice recommend weighing resident on admission/readmission, weekly for first 4 weeks after admission, and at least monthly thereafter
- Crucial that weight is obtained accurately and consistently
  - -Facility-wide scales should be calibrated and functioning appropriately
  - A consistent process for weighing residents should be in place

#### Weights - Consistent Process

Weigh the resident:

- · Wearing a similar type of clothing
- At approximately the same time of the day (preferably before breakfast)
- Using the same scale
- Either consistently wearing or not wearing orthotics or prostheses



88


#### Accurate Weights - Importance


- Weight is a parameter that reflect resident's nutritional status
- Fluid loss or retention can cause short-term weight changes; abrupt weight changes along with change in food intake are signs of fluid and electrolyte imbalance
- Weight loss may be important indicator of a change in resident's health status or sign of malnourishment
- If significant weight loss noted, health care team reviews for possible causes


89

#### Weight Component- Balance Beam

- Has 2 poise bars upper and lower
- May include pounds only or pounds on the top part of each bar and kilograms on the bottom part of each bar; may have interchangeable pound and kilogram bars







# Weight Indicators Both upper and lower bars have movable weight indicators The weight indicator for the lower bar fits into the groove as weight is obtained Weight Indicator Weight Indicator Weight Indicator Weight Indicator Weight Indicator Weight Indicator

#### **Balance Bar and Balance Window**

- Ensure balance bar floats freely and in center of window when poise bars are set on 0 (zero) and resident is not standing on scale platform
- If balance bar is off-center and/or touching window, do not weigh resident and notify nurse



94

#### Reading the Weight

100 pounds + 38 pounds = 138 pounds

To determine weight, add the value for the lower bar to the value for the upper bar.



NCDHHS/DHSR/HCPEC | Module AA Measurement | July 20

#### 95

#### Measuring Intake and Output (I&O)

- Intake (also called input) the amount of fluid taken in by the body
- Output the amount of fluid lost from the body
- Intake and output are typically seen together and commonly abbreviated I&O
- For fluid balance to occur fluid intake roughly equals fluid output

#### **Intake and Output**

- Ordered by the doctor, found on the care plan, and by the directive from the nurse
- Typically calculated at end of each shift and totaled every 24-hours
- · Documented on a facility-specific form
- · Calculations and totals based on the milliliter (mL)



97

#### Milliliter (mL)

- · A unit of measure in the metric system
- Fluids measured using the milliliter (mL)
- Another unit of measure used in health care is called the cubic centimeter (cc); should use mL instead of cc when documenting fluids
- 1 milliliter is equal to 1 cubic centimeter
- Most people familiar with the teaspoon; there are 5 mL or 5 cc in a teaspoon


98


#### **Graduate**


- Accurate measuring device for fluids when resident is on I&O
- Fluid for I&O is measured and documented in milliliters (mL)
- · Measure fluid at eye-level on a flat surface
- If both intake and output to be measured with the graduate, two separate graduates used and labeled



# Units of Measure for the Graduate The cc (cubic centimeter) side of the measurement scale is used in health care Recall 1 cc on the graduate = 1 milliliter (mL) Shortest line represents either 25 cc/mL or 75 cc/mL Longest lines represent multiples of 50 cc/mL







103

#### **Determining Oral Fluids as Intake**

- To determine intake, nurse aide must know serving sizes of facility specific containers
- Two methods
  - -Measures
  - -Using fractions
- Follow facility policy and/or procedure when determining intake of fluids during and between meals

104

#### **Typical Serving Sizes of Liquids**

- Water glass = 240 mL
- Tea glass = 180 mL
- Juice glass = 120 mL
- Milk carton = 240 mL
- Coffee cup = 240 mL
- Soft drink can = 360 mL
- Gelatin = 120 mL
- Soup bowl = 180 mL



#### **Determining Intake - Example**

- The glass with the resident's apple juice holds 240 mL of fluid
- Using the graduate and measuring apple juice left in glass, the resident did not drink 120 mL of apple iuice
- Subtracting 120 mL of apple juice the resident didn't drink from number of mL the glass holds
- · Resident drank 120 mL of apple juice

106

#### **Determining Intake Using Fractions**

- To determine fluid intake using fractions, the nurse aide needs a basic understanding of fractions in relation to a whole, which is 1
- The entire bar = 1 and includes
  - -4 equal parts or 4 fourths
  - -2 equal parts or 2 halves
  - -3 equal parts or 3 thirds


WHOLE

1/4 1/4 1/4 1/4

1/2 1/2

1/3 1/3 1/3

107



#### **Determining Intake Using Fractions**

- 240 mL in the resident's coffee cup
- Resident drank 2/3 cup of coffee
- Resident drank 160 mL of coffee

$$\frac{2}{3}$$
 of 240 mL of coffee

$$\frac{2}{3}$$
 of 240 mL of coffee  
 $\frac{2}{3}$   $\times \frac{240}{1}$  =  $\frac{160}{1}$  = 160 mL

109


#### **Determining Intake Using Fractions**

- 240 mL in the resident's coffee cup
- Resident drank 1/2 cup of coffee
- Resident drank 120 mL of coffee

$$\frac{1}{2}$$
 of 240 mL of coffee

$$\frac{1}{2} \times \frac{240}{1} = \frac{120}{1} = 120 \text{ mL}$$

110



#### **Determining Intake Using Quickly Fractions**

- Determine what 1 part of total coffee equals by dividing the total mL amount of coffee (240) by 3 = 80
- Resident drank 2 of the 3 parts of the coffee, so multiply 80 (which is 1 part) by 2 because the resident drank 2 of the 3 parts = 160
- The resident drank 160 mL of the coffee

 $\frac{2}{3}$  of 240 mL of coffee

112

#### **Milliliters and Ounces**

- A common conversion in health care is changing (or converting) ounces to milliliters
- 1 ounce = 30 mL
- To convert ounces to milliliters, simply multiply number of ounces by 30

113

#### Conversion Problem - Apple Juice

- 8 fluid ounces of apple juice in the glass
- 1 fluid ounce = 30 milliliters (mL)
- 8 x 30 milliliters (mL) = 240 milliliters (mL)
- 240 mL of apple juice in the glass



#### **Milliliters and Cubic Centimeters**

- Devices in health care are often marked in cubic centimeters (cc)
- A common conversion is changing (or converting) cc to milliliters (mL)
- 1 cubic centimeter (cc) = 1 milliliter (mL)
- To convert cc to mL, simply use the exact number measured

115

#### **Conversion Problem – Grape Juice**

- 120 cubic centimeters (cc) of grape juice in the glass
- 1 cc = 1 milliliter (mL)
- 120 mL of grape juice in the glass




116


#### Fluids Considered as Output

- Urine
- Vomitus
- Diarrhea
- Wound drainage
- Gastric suction material





118



119

#### Urinal

- A plastic, elongated device used by individuals for collection of expelled urine
- Meant for single-resident use
- As a measuring device, marked in ounces and cc (same as mL), with 100 cc (mL) increments marked lines and 50 cc (mL) unmarked lines between



#### Commode Hat (Specimen Pan)

- Plastic collection container placed under commode lid
- Used when resident on bathroom privileges and on measured output
- · Used to obtain urine or stool specimen
- Marked in ounces and cc (same as mL); grooved edge

T HANDA

NOT FRANCE

121

#### **Emesis Basin**


- A plastic, shallow basin shaped like a kidney that fits against resident's neck and collects bodily fluids
- · Used during mouthcare
- If resident is nauseated, used to collect vomitus
- Marked in ounces and cc (same as mL) with 100 cc (mL) increments


122

#### Measuring Urine Example #1

- Long line = 200 cc (mL)
- Short line = 25 cc (mL)
- Amount = 225 cc (mL)
- Document = 225 mL









# Identifying Food Intake • Nurse aide compares amount of food eaten with amount of food served • To measure food intake, nurse aide needs a basic understanding of percentages in relation to a whole, which is 100%

25%

25%

127

#### **Understanding Percentages of a Whole**

- The entire circle = 100% and includes 4 equal parts
- Each 25% is 1/4 of the whole
- 25% + 25% = 50% (or  $\frac{1}{2}$  of the whole)
- 25% + 25% + 25% = 75% (or  $\frac{3}{4}$  of the whole)




128

#### **Identifying and Documenting Food Intake**


Follow facility procedure regarding identifying and documenting food intake, for example

- Listing exact food eaten (all of chicken, all of green beans, ½
   of mashed potatoes, ¼ of biscuit, all of pie)
- Using specific words (all, good, fair, poor, refused)
- Using percentages of food eaten (100%, 75%, 50%, 25%, 0%)








130




131









The End					
NORTHE (DUCCHE COMPANY) AND					